



Dow AgroSciences

GALLANT* SUPER

Herbicide

**Post-Emergence control of grasses
in Broad Leaf and Tree crops**

Get more with
GALLANT* SUPER

WEEDS



Cynodon Dactylon



Rye Grass



Brome Grass




Digitaria Scalarum

CROPS



Tomatoes



Peas



Cotton



Potatoes



GALLANT* SUPER

Active Ingredient: Haloxyfop - R methyl Ester - 108g/Lt

FEATURES & BENEFITS:

- Gallant Super has minimal environmental impact as it rapidly breaks down in the soil and water by microbial action.
- Gallant Super is quickly absorbed through plant cuticle becoming rainfast within an hour after application.
- With Gallant, there is no need to disturb previous crop residue, this conserves soil moisture, reduces erosion and entails a low energy output for cultivation.
- Active in most climatic conditions. As long as grass weeds are growing, Gallant Super will give good control whatever the weather conditions.

TIMING OF APPLICATION:

CROP: Grass weeds may be treated at any time provided the pre harvest intervals are observed.

WEEDS: Grass Weeds must be actively growing and not under stress during application as this may limit uptake and translocation.

For annual grasses, best results are obtained at 2-6 leaf stage, while with Paspalum Spp, best results are obtained at early flowering stage.

APPLICATION RATES:

Grass weeds that will normally be controlled

| * Type of Grass | Litres/Ha | 20 Litres Knasack | Timing |
|--|------------|-------------------|--|
| Annual Grasses | | | |
| Wild oats - <i>Avena fatua</i> | 0.5 Lt | 50ml | Always apply when grass is actively growing 2-3 leaf to early tillering and no more than 12cm high. |
| Brome grass - <i>Bromus spp</i> | 0.5 Lt | | |
| Rye grass - <i>Lolium temulentum</i> | 0.5 Lt | | |
| Love grass - <i>Setaria Verticillata</i> | 0.5 Lt | | |
| False star grass - <i>chloris rynchnothrix</i> | 0.5 Lt | | |
| Crows foot grass - <i>Eleusine Indica</i> | 0.5 Lt | | |
| Crab finger grass - <i>Digitaria sanguinalis</i> | 0.5 Lt | | |
| Guinea-fowl grass - <i>Rottbelia aestivum</i> | 0.6 Lt | 60ml | |
| Volunteer wheat - <i>Triticum aestivum</i> | 1.0 Lt | 100ml | |
| Perennial Grasses | | | |
| Buffalo grass - <i>Panicum maximum</i> | 1.0 Lt | 100ml | Apply when grass is 15-30cm high. A further application may be necessary if regrowth occurs from rhizomes or new seed. |
| - <i>Panicum trichocladum</i> | 1.0 Lt | 100ml | |
| Blue couch - <i>Digitaria scalarum</i> | 1.2 Lt | 120ml | |
| Kikuyu grass - <i>Pennisetum cladeatum</i> | 1.2 Lt | 120ml | |
| Sudan grass - <i>Sorghum halepense</i> | 1.2 Lt | 110ml | |
| Star grass - <i>Cynodon dactylop</i> | 1.0 - 2 Lt | 100-200ml | Increase rate in dry conditions to 2.0 - 2.4Lt/ha |
| Sword grass - <i>Imperata cylindrical</i> | 1.0 - 2 Lt | 100-200ml | |

READ THE LABEL BEFORE USE



Lachlan@griculture

Marketed & Distributed by: **LACHLAN KENYA LIMITED**. Tel: +254 2073912/3/4
 +254 0722 209 474 Fax: +254 2060260 . P.O. BOX 49470, 00100 Nairobi.
 Email: lachlan@griculture.co.ke

*** Trademark of Dow Agrosciences**