

NEEMROC E.C. 0.03% contains alcoholic extract of neem seeds, leaves in drip irrigation systems. Neemroc E.C. is used to control Diamond backmoth & thrips on brassicas.

DIRECTIONS FOR USE:

Shake well before use.

RATES OF APPLICATION:

2.5mls/1Lt.

Compatibility: Neemroc E.C is compatible with many pesticides.

Precautions: It should be handled carefully to avoid contamination. It is not to be taken orally. Avoid contamination of eyes and body, eat or drink while handling this product.

Always adhere to the following instructions when handling this product.

- Use protective clothing, gloves, gumboots headscarf or appropriate hat and face shield.
- Avoid breathing spray mist. Never spray against the wind.
- Avoid direct contact with skin and eyes.
- Change used clothing and bathe thoroughly with enough water and soap.
- Thoroughly clean spraying equipment and working attire.

Storage: Store in original packaging in a cool, dry, well ventilated area. Store away from sun, under lock and key. Keep away from food, drink and foodstuffs.

Product and container disposal: Puncture empty containers. Do not re-use for any other purpose. Bury the pack away from human habitation and in an area where contamination will not occur.

Leakage or spillage: Should this occur, soak up materials with sand, soil or saw dust. Collect the contaminated materials and bury them in a safe place. Wash the contaminated areas thoroughly.

FIRST AID INSTRUCTIONS: If poisoning is suspected, stop work immediately. Check for any signs of distress. Remove clothing if contaminated and get out of area. If in eyes, rinse immediately with plenty of water for at least ten minutes. In case of contact with skin, wash with water and soap. If inhaled, move the patient to fresh air and obtain medical attention. If swallowed, induce vomiting by gently touching the back of the throat; never induce vomiting if patient is unconscious.

TOXICOLOGICAL INFORMATION

Symptoms of Poisoning: Mild irritation of the eyes, nose, throat, drowsiness, dizziness, headache and dry cracked skin.

Antidote: No specific Antidote.

Note to Physician: Treat symptomatically.

Notice to users: This pest Control Product is to be used only in is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use or store a Pest Control Product under unsafe conditions.

WARRANTY: Seller's guarantee is limited to the terms set out on the label and the subject handling of this product and accepts the product in that condition.

**READ THE LABEL BEFORE USING
(SOMA KIBANDIKO KABLA YA KUTUMIA DAWA)**

NEEMROC E.C. 0.03% w/w

An extract of herbaceous plants used to control insects in Brassicas.

REGISTRATION NO. PCPB (TCR) 0049

(Nambari ya Usajili)

GUARANTEE (DHAMANA): 0.03% Azadirachtins

Date of Manufacture (Tarehe ya Kuindwa)

Batch No. (Nambari ya furushi)

SHELF LIFE:

One year from date of manufacture if stored in a cool, dark, dry well ventilated place in original sealed containers.

MUDA WA KUDUMU

Mwaka moja kutoka tarehe ya kutengenezwa ikiwa kwa mkoba uliofunikwa vizuri na kuhifadhiwa pasipo na joto nyingi wala unyevu.

Manufactured by:

(Watengenezaji)

Saroneem Biopesticide Limited
Baba Dogo, Baba Dogo House
P.O. Box 62373 - 00620,
Nairobi, Kenya
Tel: 020 8562675

Distributed by:

(Wasambazaji)

Lachlan Kenya Limited
P.O. Box 49470,0010
Tel: 020 2073912/3/4
Mobile: 0722 209474
Fax: 020 2060260

**NET CONTENT..... LITRES
(KIASI)**



Tingisha vizuri kabla ya kutumia NEEMROC+ 0.03% + Azadiractin ina mfinyo wa mbegu za moarobaine, matawi na miti mingine ya kienejeji. Bidhaa hii imetengenezwa maalum kunyunyiziwa kwa njia ya kutonesha

JINSI YA KUFANYA KAZI

Neemroc E.C. inatumika kuzuia wadudu kwa mchanga haswa.

MAAGIZO YA KUTUMIA

Pasipo maelezo mengine, Neemroc E.C yastahili kuongezwa kwa maji ya kunyunyizia kwa njia ya kutonesha katika kipimo cha ½ lita kwa hectare moja kila wiki.

TAHADHARI YA KUTUMIA

Hii ni dawamu ya kuwanganamiza wadudu, iliyotengenezwa kutokana na utaalumu wa kibiolajia na isiyo na sumu nyingi. Hata hivyo, lazima itumwe vyema kuzuia maradhi yeyote. Haiatahili unywe, chungu isingie macho au mwili usivute sigara, kula au kunywa unapoitumia.

Fuata maelezo kila wakati unapotumia dawa hii:

- Vaa mavazi yanayoweza kuingia ili dawa isiguzwe mwili wako kama vile: aproni, viziuzi vya mikono, na viatu vya gumboots
- Nyunyiza dawa ukiwa upande upepo unapotoka kuzuia kuvuta harufu ya dawa.
- Chungu dawa isiguzwe ngozi na macho.
- Badilisha nguo, oga na maji safi na sabuni baada ya kunyunyiza dawa.
- Osha vyombo vya kunyunyizia kabisa pamoja na nguo baada ya kazi.

KUHIFADHI: Hifadhi kwenye mkoba uliyo nunulia dawa pahali pakavu penye baridi kiasi. Weka mbali na jua, weka mbali na chakula na vinywaji.

UTUPAJI: Osha mikoba uliyo tumia mara tatu. Usitumie tena kwa njia yeyote mkoba mtupu wa dawa hii haribu na utupe kwenye shimo refu.

KUMWAGIKA/KUVUNJA: Hii ina potokea, mwagia mchanga, vumbi au changarawe. Okota hiyo taka uitupe kwa shimo kisha funika vyema na mchanga. Osha sehemu zilizomwagikiwa sawasawa.

MAZINGARA NA WANYAMA WA PORI

Inadhuru samaki, Haidhuru nyuki, Usichafue mabwawa, mito au mikoba mitupu.

HUDUMA YA KWANZA: Dalili ya sumu inaposhukiwa wacha kazi haraka, toa nguo zilizoguzana na uondohe mahali hapo. Kama macho yameguzwa osha vizuri na maji kwa muda wa dakika kumi, kama ngozi imeguzwa oga na sabuni na maji, kama imepumuliwa weka mgonjwa mahali pana hewa safi, ikimezwa tapisha mgonjwa kwa kushika koo, usimtapishe anapozirai.

DALILI YA SUMU: kuumwa na kichwa, kuchafuka roho n.k.